



Hip Evaluation Report

Owner Copy

Reference #: 902901
Practice #: 65468 SAWYER

Report Date: 9/24/2012
Radiography Date: 9/18/2012
Date Received: 9/24/2012

Owner:
STEFANI WHITEMAN

PennHIP Member:
DR. SUSAN M. NEWELL
OCEAN STATE VETERINARY SPECIALISTS
1480 COUNTY TRAIL
EAST GREENWICH, RI 02818
UNITED STATES

ANIMAL			
OCEAN STATE'S TOM SAWYER (SAWYER)		Reg. #:	
CANINE / LABRADOODLE CROSS		Microchip: 985121012093440	
Date of Birth: 3/7/2012	Sex: M	Weight: 22 lbs.	Age: 6 mo.
		Tattoo:	
RESULTS			
LEFT	Distraction Index (DI)	0.34	DI is greater than 0.30 with no radiographic evidence of DJD. There is an increasing risk of developing DJD as the DI increases; low risk when DI is close to 0.30, high risk when DI is close to 0.70 or above.
	Degenerative Joint Disease (DJD)	None	
	Cavitation	No	
	Other Findings	Not Applicable	
RIGHT	Distraction Index (DI)	0.34	DI is greater than 0.30 with no radiographic evidence of DJD. There is an increasing risk of developing DJD as the DI increases; low risk when DI is close to 0.30, high risk when DI is close to 0.70 or above.
	Degenerative Joint Disease (DJD)	None	
	Cavitation	No	
	Other Findings	Not Applicable	

Please note that the PennHIP DI is a measure of hip joint laxity, it does not allude to a "passing" or "failing" hip score.

LAXITY PROFILE RANKING										
The laxity profile ranking is based on the hip with the greater laxity (DI). This interpretation is based on a cross-section of 2,326 CANINE animals of the LABRADOODLE CROSS breed. The median DI for this group is 0.52.										
Percentiles										
	90th	80th	70th	60th	50th	40th	30th	20th	10th	
> 90th					Median					< 10th
↑										
The chart above indicates the ranking of your animal's passive hip laxity (DI) in relation to all CANINE animals of the LABRADOODLE CROSS breed in our database. This result means that 1) your animal's hips are tighter than over 90% of the animals in this group, and 2) your animal's hip laxity is in the tighter half of the laxity profile. Breed-specific evaluations are analyzed semi-annually. Consequently, the average laxity and range of laxity for any given group will change over time.										

PennHIP does not make specific breeding recommendations. Selection of sire and dam for mating is the decision of the breeder.

NOTE: As a minimum breeding criterion, we propose that breeding stock be selected from the population of animals having hip laxity in the tighter half of the breed (to the left of the median mark on the graph). Higher selection pressure equates to more rapid expected genetic change per generation.

By implementing selection based on passive hip laxity, we expect the breed average DI over the years to move toward tighter hip configuration, meaning lower hip dysplasia susceptibility. The PennHIP database permits scientific adjustment of criteria to reflect these shifts; the average laxity and range of laxity for a particular breed will change over time.

PennHIP / Veterinary School of the University of Pennsylvania / 3800 Spruce Street / Philadelphia, PA 19104

www.pennhip.org