



Hip Evaluation Report

Owner Copy

Reference #: 898606
Practice #: 7724

Report Date: 2/7/2012
Radiography Date: 1/3/2012
Date Received: 2/3/2012

Owner:
DEBORAH LEE


PennHIP Member:
DR. JAMES R. MILES
ST. CHARLES VETERINARY CLINIC
530 DUNHAM RD
ST CHARLES, IL 60174
UNITED STATES

ANIMAL	
AUSSI DOG'S CHOCOLATE KISS (MOCHA)	Reg. #:
CANINE / LABRADOODLE CROSS	Microchip: 014 074 530
Date of Birth: 2/4/2011 Sex: F Weight: 23 lbs. Age: 11 mo.	Tattoo:

RESULTS			
LEFT	Distraction Index (DI)	0.41	DI is greater than 0.30 with no radiographic evidence of DJD. There is an increasing risk of developing DJD as the DI increases; low risk when DI is close to 0.30, high risk when DI is close to 0.70 or above.
	Degenerative Joint Disease (DJD)	None	
	Cavitation	No	
	Other Findings	Not Applicable	
RIGHT	Distraction Index (DI)	0.42	DI is greater than 0.30 with no radiographic evidence of DJD. There is an increasing risk of developing DJD as the DI increases; low risk when DI is close to 0.30, high risk when DI is close to 0.70 or above.
	Degenerative Joint Disease (DJD)	None	
	Cavitation	No	
	Other Findings	Not Applicable	

Please note that the PennHIP DI is a measure of hip joint laxity, it does not allude to a "passing" or "failing" hip score.

LAXITY PROFILE RANKING
The laxity profile ranking is based on the hip with the greater laxity (DI). This interpretation is based on a cross-section of 2,077 CANINE animals of the LABRADOODLE CROSS breed. The median DI for this group is 0.53.



The chart above indicates the ranking of your animal's passive hip laxity (DI) in relation to all CANINE animals of the LABRADOODLE CROSS breed in our database. This result means that 1) your animal's hips are tighter than approximately 80% of this group of animals (alternatively, 20% of the group has tighter hips than your animal), and 2) your animal's hip laxity is in the tighter half of the laxity profile. Breed-specific evaluations are analyzed semi-annually. Consequently, the average laxity and range of laxity for any given group will change over time.

PennHIP does not make specific breeding recommendations. Selection of sire and dam for mating is the decision of the breeder.

NOTE: As a minimum breeding criterion, we propose that breeding stock be selected from the population of animals having hip laxity in the tighter half of the breed (to the left of the median mark on the graph). Higher selection pressure equates to more rapid expected genetic change per generation.

By implementing selection based on passive hip laxity, we expect the breed average DI over the years to move toward tighter hip configuration, meaning lower hip dysplasia susceptibility. The PennHIP database permits scientific adjustment of criteria to reflect these shifts; the average laxity and range of laxity for a particular breed will change over time.

PennHIP / Veterinary School of the University of Pennsylvania / 3800 Spruce Street / Philadelphia, PA 19104

www.pennhip.org